



KONGERIKET NORGE
The Kingdom of Norway

REC'D 26 JUL 2004

WIPO PCT

Bekreftelse på patentsøknad nr
Certification of patent application no

20033078

▷ Det bekreftes herved at vedheftede dokument er nøyaktig utskrift/kopi av ovennevnte søknad, som opprinnelig inngitt 2003.07.04

▷ It is hereby certified that the annexed document is a true copy of the above-mentioned application, as originally filed on 2003.07.04

2004.07.13

Line Reum

Line Reum
Saksbehandler

**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)



552 5350

2003 -07-04

**PATENTSTYRET**
Styret for det industrielle rettsvernADDRESSE
Postboks 6140 Dn
Kobenhavngaten 10
0033 OsloTELEFON
22 34 72 00
TELEFAKS
22 39 73 01BANKGIRO
8276 01 00192
FORETAKSNUMMER
971520157

PATENTSTYRET

Søknad om patent

03-07-04*20033078

Skal utfylles av Patentstyret

Behandler medlem **KL**
Int. Cl. **A61K****Alm. tilgj. 05 JAN 2005**Søknaden skal være revideres
(for og etter innlegg)JOH/vv
Case 11Oppfinnelsens
betegnelse**Fish-protein hydrolysate***Hydrolysat av fiskeprotein*Hvis søknaden er
en internasjonal søknad
som videreføres etter
patentlovens § 31:

Den internasjonale søknads nummer

Den internasjonale søknads inngivelsesdag

Søker

Søker skal angi adresse.
Hvis patent anses av færet
opplysning om hvem som skal
være ansvarlig for å motta
meddelelser fra Patentstyret på
vegne av søkeren.**Thia Medica AS**
c/o siv. øk. Terje Moe
Kalfarveien 57 A
5018 Bergen

(For søker som har verd, på neste side)

Søker er en enkeltperson eller en småbedrift, eller flere søke i fellesskap med fast ansatte som til-
sammen utfører 20 årsverk eller mindre (på søknadstidspunktet). Det er søkers ansvar å krysse av her
for å oppnå laveste satser for søknadsavgift. NB! se også utfyllende forklaring på siste side.

Oppfinner.

Navn (et, et eller flere)
(For søker som har verd, på neste side)**Rolf Berge**
Tjørnhaugen 50
5152 Bønes

Fu mektig

AS BERGEN PATENTKONTOR
Strandgaten 198, 5004 BergenHvis søknad tidligere
er inngitt, eller
utenfor riket.

(For søker som har verd, på neste side)

Prioritet kreves fra dato sted nr.
Prioritet kreves fra dato sted nr.
Prioritet kreves fra dato sted nr.

Hvis syde t søknad:

Den opprinnelige søknads nr.: og deres inngivelsesdag

Hvis Ltskilt søknad:

Den opprinnelige søknads nr.: begjært inngivelsesdag

Deponert kultur av
mikroorganisme:

Søknaden omfatter kultur av mikroorganisme. Oppgi også deponeringssted og nr.

Utlevering av prøve av
kulturen:Prøve av den deponerte kultur av mikroorganisme skal bare utleveres til en særlig sakkyndig.
jfr patentlovens § 22 åttende ledd og patentforskriftens § 38 første leddAngivelse av tegnings-
figurer som ønskes
publisert sammen med
søkmendraget

Fig. nr.

1b

PATENTSTYRET

03-07-04*20033078

FIELD OF INVENTION

The present invention relates to the use of a fish protein hydrolyzate (FPH). The FPH material lowers the concentration of cholesterol in plasma, and triglycerides in the liver.

5 FPH also induces a favourable change in the fatty acid pattern, and lowers the concentration of homocysteine in plasma. A preferable embodiment of the invention relates to the use of FPH as an anti-atherogenic and cardio protective agent, either given as a pharmaceutical or as a functional food.

10

BACKGROUND OF THE INVENTION

The fish farming industry has grown enormously both in Norway and world-wide during the recent years, especially Salmon. Much of the fish are sold to the consumer as gutted,

15 whole fish, but significant amounts are sold as fillets. Only 50-70 % of the salmon is fillets, whereas the rest is sold as low-valued products such as fish meal and fish ensilage.

Through enzymatic treatment the fish meat and also the fish frames can be separated

20 into an aqueous fraction rich in proteins, called fish protein hydrolysate (FPH). The enzymatic hydrolysing process is highly controllable, and the products are reproducible and well defined.

Surprisingly, the present inventors have shown that a fish protein hydrolysate in accordance with the invention has several beneficial biological effects, and that such a material can be used as a pharmaceutical.

- 5 We have shown that the fish protein hydrolysate lowers the concentration of plasma cholesterol and homocysteine, and also lowers the concentration of hepatic triacylglycerols. Based on these findings, it is anticipated that the fish protein hydrolysate will have a preventive and/or therapeutic effect on stenosis, atherosclerosis, coronary heart disease, thrombosis, myocardial infarction, stroke and fatty liver.
- 10 Treatment with a fish protein material represents a new way to treat these diseases.

- The fish protein hydrolysate is especially useful as a functional protein in food products, particularly when used as a substitute for natural plasma in animal feeds and in pet foods. When used in pet foods, additional ingredients may be added to the product such
- 15 as fats, sugars, salt, flavourings, minerals, etc. The product may then be formed into chunks resembling natural meat chunks in appearance and texture. The product of the invention has the further advantages that this is readily formulated to contain necessary nutrients, is easily digested by the animals and is palatable to the animals.

20 DETAILED DESCRIPTION OF THE INVENTION

- The present invention relates to a fish protein hydrolysate for the preparation of a pharmaceutical or nutritional preparation for the treatment and/or prevention of atherosclerosis, coronary heart disease, stenosis, thrombosis, myocardial infarction,
- 25 stroke and fatty liver.

- The experimental data clearly shows that the fish protein hydrolysate according to the invention lowers the concentration of homocysteine in plasma. Homocysteine is a risk factor in diseases such as atherosclerosis, coronary heart disease, stenosis, thrombosis,
- 30 myocardial infarction and stroke, and it is thus anticipated the FPH material of the invention will be effective in preventing and treating of these diseases.

The data also shows that the level of triacylglycerols in the liver is decreased by administration of FPH, and it is anticipated that the FPH material can be used for the treatment and prevention of fatty liver.

5 A further embodiment of the present invention relates to a fish protein hydrolysate for the preparation of a pharmaceutical or nutritional composition for the treatment and/or prevention of hypercholesterolemia, as we have shown that said material is capable of lowering the plasma concentration of cholesterol.

10 A still further embodiment relates to the use of a fish protein hydrolysate for the preparation of a pharmaceutical or nutritional composition for lowering the concentration of homocysteine in the plasma. A hyperhomocysteine level can be established before the above indicated diseases are manifested. The administration of the FPH material has a general homocysteine lowering effect, and the material of the
15 present invention is thus especially suited for preventing the onset of, and lowering the risk for the above indicated diseases.

The results further indicate that the FPH material has general cardio and artery protective features, and we anticipate that the material can be given to decrease the risk
20 for artery and cardio related diseases.

An object of the present invention is to administer the material either as a prophylactic or pharmaceutical drug, or as a functional feed or food material. The material can be given to human and non-human animals.

25

A preferred embodiment of the invention relates to a feed material comprising the fish protein hydrolysate. The material can be used for feeding agricultural animal, such as gallinaceous birds, bovine, ovine, caprine or porcine mammals, domestic or pet animal, such as dog or cat, and fish or shellfish, such as salmon, cod, Tilapia, clams, oysters,
30 lobster or crabs.

A preferred embodiment of the invention uses FPH material produced by an enzymatic treatment of fish meat or frames. Preferable the enzyme composition ProtamexTM is used, and the fish is preferable salmon.

5 FIGURE LEGENDS

Figure 1 shows that the fish protein hydrolysate (FPH) decreases the concentration of cholesterol in plasma.

10 Figure 2 shows that the fish protein hydrolysate (FPH) decreases the concentration of cholesterol in the liver.

Figure 3 shows that the fish protein hydrolysate inhibits the ACAT enzyme.

15 Figure 4 shows that the fish protein hydrolysate increases the mitochondrial β -oxidation.

DEFINITIONS USED IN THE APPLICATION

20

Animals

In this context the term "animals" include mammals such as humans and farm (agricultural) animals, especially the animals of economic importance such as gallinaceous birds, bovine, ovine caprine and porcine mammals, especially those that
25 produce products suitable for the human consumption, such as meat, eggs and milk. Further, the term is intended to include fish and shellfish, such as salmon, cod, Tilapia, clams and oysters. The term also includes domestic animals such as dogs and cats.

Treatment

30

In relation to the pharmaceutical applications of the invention the term "treatment" refers to a reduction of the severity of the disease.

Prevention

The term "prevention" refers to the preventing of a given disease, i.e. a compound of the present invention is administered prior to the onset of the condition. This means that the compounds of the present invention can be used as prophylactic agents or as ingredients
5 in functional foods or feed in order to prevent the risk or onset of a given disease.

Fish protein hydrolysate

Fish protein hydrolysate is an enzymatic treated fish material. The FPH material contains high proportions of proteins.
10

ADMINISTRATION OF THE COMPOUNDS OF THE PRESENT INVENTION

As a pharmaceutical medicament the compounds of the present invention may be
15 administered directly to the animal by any suitable technique, including parenterally, intranasally, orally, or by absorption through the skin. They can be administered locally or systemically. The specific route of administration of each agent will depend, e.g., on the medical history of the animal.

20 Examples of parenteral administration include subcutaneous, intramuscular, intravenous, intraarterial, and intraperitoneal administration

If given continuously, the compounds of the present invention are each typically administered by 1-4 injections per day or by continuous subcutaneous infusions, for
25 example, using a mini-pump. An intravenous bag solution may also be employed. The key factor in selecting an appropriate dose is the result obtained, as measured by decreases in total body weight or ratio of fat to lean mass, or by other criteria for measuring control or prevention of obesity or prevention of obesity-related conditions, as are deemed appropriate by the practitioner.

30 For parenteral administration, in one embodiment, the compounds of the present invention are formulated generally by mixing each at the desired degree of purity, in a unit dosage injectable form (solution, suspension, or emulsion), with a pharmaceutically

acceptable carrier, i.e., one that is non-toxic to recipients at the dosages and concentrations employed and is compatible with other ingredients of the formulation.

Generally, the formulations are prepared by contacting the compounds of the present invention each uniformly and intimately with liquid carriers or finely divided solid carriers or both. Then, if necessary, the product is shaped into the desired formulation. Preferably the carrier is a parenteral carrier, more preferably a solution that is isotonic with the blood of the recipient. Examples of such carrier vehicles include water, saline, Ringer's solution, and dextrose solution. Non-aqueous vehicles such as fixed oils and ethyl oleate are also useful herein, as well as liposomes.

The carrier may suitably contain minor amounts of additives such as substances that enhance isotonicity and chemical stability. Such materials are non-toxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, succinate, acetic acid, and other organic acids or their salts; antioxidants such as ascorbic acid; immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids, such as glycine, glutamic acid, aspartic acid, or arginine; monosaccharides, disaccharides, and other carbohydrates including cellulose or its derivatives, glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; counterions such as sodium; and/or non-ionic surfactants such as polysorbates, poloxamers, or PEG.

For oral pharmacological compositions such carrier material as, for example, water, gelatine, gums, lactose, starches, magnesium-stearate, talc, oils, polyalkene glycol, petroleum jelly and the like may be used. Such pharmaceutical preparation may be in unit dosage form and may additionally contain other therapeutically valuable substances or conventional pharmaceutical adjuvants such as preservatives, stabilising agents, emulsifiers, buffers and the like. The pharmaceutical preparations may be in conventional liquid forms such as tablets, capsules, dragees, ampoules and the like, conventional dosage forms, such as dry ampulles, and as suppositories and the like.

In addition, the compounds of the present invention are appropriately administered in combination with other treatments for combating or preventing a specific disease.

The invention will be more fully understood by reference to the following examples. They should not, however, be construed as limiting the scope of the invention.

- 5 As a nutritional composition the FPH may be formulated in any conventional way to a feed or food product.

EXPERIMENTAL SECTION

10

The following non-limiting examples serve to further illustrate the invention.

Chemicals

- 15 [1-¹⁴C] palmitoyl-L-carnitine (54 Ci/mmol) was purchased from Amersham. The chemicals used for real-time RT-PCR was from Applied Biosystems. All other chemicals were obtained from common commercial sources and were of reagent grade.

20 Fish protein hydrolyzate (FPH)

- FPH was produced from fish flesh remnants on salmon bone frames after filleting as described previously (23) (24), the Supro 530 EX soy protein was from duPont Protein Technologies (St.Louis, MO, USA), and bovine casein sodium salt, C-8654, was from
25 Sigma-Aldrich.

Animals and treatments

- 4-5 weeks old male obese Zucker rats, Crl:(ZUC)/faBR from Charles River, Germany,
30 averaging 120 ± 3 g at the start of the experiment, were kept in a room maintained at 12 hours light-dark cycles, at a temperature of $20 \pm 3^\circ\text{C}$, and relative humidity of $65 \pm 15\%$. The day after arrival the rats were randomised and placed separately in metabolic cages and divided into three experimental groups, each of six animals. The rats were

adapted to the experimental conditions and experimental diets for 4 days, after which the faeces were collected for 7 days. The semipurified diets (Table 1), contained 20% crude protein ($N \times 6,25$) in the form of FPH or casein (control).

TABLE 1

Composition of the experimental diets

<u>g/kg diet</u>	FPH	Casein
Protein	233,9	217,6
Soybean oil ¹	100	100
Sucrose	110	110
Vitamins ²	10	10
Minerals ³	30	30
Cellulose	20	20
NaCl	-	21,8
Dextrin	496,1	490,6

¹ Fatty acid composition of the soybean oil (g/100g fat): 18:2n-6 (54.1 ± 0.5), 18:1n-9 (21.8 ± 0.2), 16:0 (11.2 ± 0.1), 18:3n-3 (6.1 ± 0.2), 18:0 (3.7 ± 0.1), 18:1n-7 (1.5 ± 0.1), 20:0 (0.5 ± 0.1), 22:0 (0.5 ± 0.1).

² Vitamins (mg/kg diet): 8 mg vit.A (4000 I.U.), 2 mg vit.D3 (1000 I.U.), 60 mg vit.E (30 I.U.), 0.1 mg vit.K (0.05 I.U.), 1000 mg choline hydrogentartrate, 4 mg thiamine, 3 mg riboflavin, 6 mg pyridoxine, 20 mg niacin, 8 mg Ca-pantothenat, 1 mg folin, 5 mg vit.B12 (0.05 I.U.).

³ Minerals (g/kg diet): 8.5 g $CaCO_3$, 6.2 g $CaHPO_4 \cdot 2H_2O$, 12.3 g KH_2PO_4 , 1.4 g $MgCO_3$, 0.4 $NaCO_3$, 0.8 g NaCl, 0.02 g $CuSO_4 \cdot 5H_2O$, 0.002 g NaF, 0.0002 g KI, 0.2 g $FeSO_4 \cdot xH_2O$, 0.05 g $ZnSO_4 \cdot xH_2O$.

The animals were daily offered equal feed rations, which were adjusted to meet the demand of the growing animal. The animals had free access to tap water. The rats were fed for 22 or 23 days after acclimatisation (three rats from each group were killed on day 22 and the rest on day 23), and the body weight was measured weekly. At the end of the feeding period, the animals were anaesthetised subcutaneously by 1:1 Hypnorm[®]/Dormicum[®] (Fentanyl/fluanisone-Midazolam), 0.2 mL/100 g body weight. Cardiac puncture was performed to collect blood samples (in heparin), and the liver was dissected. Parts of the liver were immediately frozen in liquid N₂, while the rest of the

liver was chilled on ice for homogenisation. The protocol was approved by the Norwegian State Board of Biological Experiments with Living Animals.

Preparation of subcellular fractions

5

Livers from the rats were homogenised individually in ice-cold sucrose-solution (0.25 mol/L sucrose in 10 mmol/L HEPES buffer pH 7.4 and 1 mmol/L EDTA) using a Potter-Elvehjem homogeniser. The subcellular fractions were isolated as described in Berge, R. K. et al (Berge, R. K., Flatmark, T. & Osmundsen, H. (1984), Enhancement of long-chain acyl-CoA hydrolase activity in peroxisomes and mitochondria of rat liver by peroxisomal proliferators. Eur J Biochem 141: 637-644). Briefly, the homogenate was centrifuged at 1 000 x g for 10 min to separate the post-nuclear from the nuclear fraction. A mitochondrial-enriched fraction was prepared from the post-nuclear fraction at 10 000 x g for 10 min. A peroxisome-enriched fraction was prepared by centrifugation of the post-mitochondrial fraction at 23 500 x g for 30 min. A microsomal-enriched fraction was isolated from the post-peroxisomal fraction at 100 000 x g for 75 min. The remaining supernatant was collected as the cytosolic fraction. The procedure was performed at 0-4°C, and the fractions were stored at -80°C. Protein was assayed using the BioRad protein assay kit (BioRad, Hercules, CA) and bovine serum albumine as standard.

20

Enzyme assays

Carnitine palmitoyltransferase I (CPT-I) activity was measured essentially as described by Bremer (Bremer, J. (1981) The effect of fasting on the activity of liver carnitine palmitoyltransferase and its inhibition by malonyl-CoA. Biochim Biophys Acta 665: 628-631). The assay for CPT-I contained 20 mmol/L HEPES pH 7.5, 70 mmol/L KCl, 5 mmol/L KCN, 100 µmol/L palmitoyl-CoA, 10 mg BSA/mL, and 0.6 mg tissue protein/mL. The reaction was started with 200 µmol/L [methyl-¹⁴C] L-carnitine (200 cpm/nmol). Assay conditions for CPT-II were identical except that BSA was omitted and 0.01% Triton X-100 was included. Tissue protein concentration was 2.5 µg/mL. Acyl-coenzyme A cholesterol acyltransferase (ACAT) was measured by using 130 mg protein and ¹⁴C-oleyl-CoA as substrate. The product was separated on TLC plates using

30

hexane:diethylether:acetic acid (80:20:1) as the mobile phase, and counted in a scintillation counter (Win Spectral 1414 liquid scintillation counter, Wallac). 3-Hydroxy-3-methylglutaryl (HMG)-CoA reductase was measured by using 80 mg protein and ^{14}C -HMG-CoA as a substrate. The product was separated on TLC plates
5 using acetone:benzene (1:1) as the mobile phase, and counted in a scintillation counter. Fatty acid synthase was measured as described by Roncari (Roncari, D. A. (1981) Fatty acid synthase from human liver. Methods Enzymol 71 Pt C: 73-79), modified according to Skorve *et al.* (Skorve, J., al-Shurbaji, A., Asiedu, D., Bjorkhem, I., Berglund, L. & Berge, R. K. (1993) On the mechanism of the hypolipidemic effect of sulfur-substituted
10 hexadecanedioic acid (3-thiadicarboxylic acid) in normolipidemic rats. J Lipid Res 34: 1177-1185), and acetyl-CoA carboxylase was determined by measuring the amount of $\text{NaH}^{14}\text{CO}_3$ incorporated into malonyl-CoA.

Lipid analysis

15

Lipids in whole liver and heparinised plasma were measured in the Tecnicon Axon system (Miles, Tarrytown, NY), with the Bayer triglyceride and cholesterol enzymatic kits (Bayer, Terrytown, NY) and the PAP 150 phospholipid enzymatic kit (bioMérieux, Lyon, France). Liver lipids were first extracted according to Bligh and Dyer (Bligh, E.
20 G. & Dyer, W. J. (1959) A rapid method of total lipid extraction and purification. Can J Biochem Physiol 37: 911-91.

Faecal sterols

25 Faecal total bile acids were prepared according to Suckling *et al.* (Suckling, K. E., Benson, G. M., Bond, B., Gee, A., Glen, A., Haynes, C. & Jackson, B. (1991) Cholesterol lowering and bile acid excretion in the hamster with cholestyramine treatment. Atherosclerosis 89: 183-190) with some modifications. Two mL of NaBH in ethanol (mg/mL) was added to 0.1 g of powdered dry feces. The mixture was allowed to
30 react for 1 hour at ambient temperature, after which 50 μL of 2 mol/L HCl was added to remove any excess of NaBH. Neutral sterols were extracted from the samples with n-hexan (two consecutive washings) before the samples were hydrolysed over night with 200 μL 10 mol/L NaOH at 110°C. 240 μL of the hydrolysate together with 2.8 mL water

was applied to Bond Elut C¹⁸ columns (Varian, 200mg, 3 mL), that had previously been activated by 3 mL methanol and 3 mL water. Bile acids were retained in the columns, which were washed twice with 3 mL of 20% methanol in water, before the bile acids were eluted with 3 mL of methanol. The bile acids were air-dried at 45°C and resolved in 1 mL of isopropanol. Total bile acids were determined enzymatically using a total bile acid diagnostic kit (Sigma 450A) on the Technicon Axon system.

Amino acids

Amino acids in the diets were determined after hydrolysis in 6 M HCl at $110 \pm 2^\circ\text{C}$ for 22 hours and pre-derivatisation with phenylisothiocyanate according to the method of Cohen and Strydom (34). Total cysteine in the feeds was determined after oxidation of cysteine and cystine with 9:1 performic acid (88 %): H_2O_2 (30 %) (v/v) to yield cysteic acid. The samples were then hydrolysed in 6 M HCl at $110 \pm 2^\circ\text{C}$ for 22 hours and further treated as the amino acid analysis described above. Amino acids in liver and plasma were determined in a Biochrom 20 plus Amino Acid Analyzer (Amersham Pharmacia Biotech, Sweden) equipped with a lithium column with post column ninhydrin derivatization as previously described (24). Prior to analysis, liver samples were extracted and deproteinated by the addition of 2 volumes of 5% sulfosalicylic acid, kept on ice for 30 min and centrifuged at $5\,000 \times g$ for 15 min. The supernatants were mixed 4:1 (v/v) with internal standard (2.5mmol/L Norleucine in 0.1 mol/L HCl). Plasma samples were mixed 1:1 with internal standard (1mmol/L Norleucine in 0.1 mol/L HCl), centrifuged at $10\,000 \times g$ for 5 min before the supernatant was centrifuged in a filter tube (cut off 10 kDa, Biomax PB polyethersulfone membrane, Millipore Corp., USA) at $10\,000 \times g$ for 30 min.

Fatty acid composition

Fatty acids were extracted from the samples with 2:1 chloroform: methanol (v/v) (35). The samples were filtered, saponified and esterified in 12% BF_3 in methanol (v/v). Fatty acid composition of total lipids from liver and plasma was analysed using methods described by Lie and Lambertsen (Lie, O. & Lambertsen, G. (1991) Fatty acid composition of glycerophospholipids in seven tissues of cod (*Gadus morhua*),

determined by combined high-performance liquid chromatography and gas chromatography. J Chromatogr 565: 119-129). Fatty acid methyl esters were separated using a Carlo Erba gas chromatograph ('cold on column' injection, 69°C for 20 s, increase at 25°C min⁻¹ to 160°C and hold at 160°C for 28 min, increase at 25°C min⁻¹ to 190°C and hold at 190°C for 17 min, increase at 25°C min⁻¹ to 220°C and hold at 220°C for 9 min) equipped with a 50 m CP-sil 88 (Chrompack, Middelburg, The Netherlands) fused silica capillary column (i.d. 0.32 mm). The fatty acids were identified by retention time using standard mixtures of methyl esters (Nu-Chek-Prep, Elyian, MN, USA). The fatty acid composition (weight percentage) was calculated using an integrator (Turbochrom Navigator, Version 4.0) connected to the GLC.

Lipids were extracted from plasma triacylglycerol-rich lipoprotein fraction using a mixture of chloroform and methanol, and separated by thin layer chromatography on silica gel plates (0.25mm Silica gel 60, Merck) developed in hexane-diethyl ether-acetic acid (80:20:1, v/v/v) and visualized using Rhodamine 6G (0.05% in methanol, Sigma) and UV light. The spots were scraped off and transferred to tubes containing heneicosanoic acid (21:0) as internal standard. BF₃-methanol was added to the samples for transesterification. To remove neutral sterols and non-saponifiable material, extracts of fatty acyl methyl esters were heated in 0.5mol/L KOH in ethanol-water solution (9:1). Recovered fatty acids were then re-esterified using BF₃-methanol. The methyl esters were analyzed on a GC8000Top gas chromatograph (Carlo Erba Instrument), equipped with a flame ionization detector (FID), programmable temperature of vaporization injector, AS 800 autosampler (Carlo Erba Instrument) and a capillary column (60m x 0.25mm) containing a highly polar SP 2340 phase with film thickness 0.20µm (Supelco). The initial temperature was 130°C, heating 1.4°C/min to final temperature 214°C. The injector temperature was 235°C. The detector temperature was 235°C, using hydrogen (25mL/min), air (350 mL/min) and nitrogen as make-up gas (30mL/min). The samples were run with constant flow using hydrogen as a carrier gas (1.6 mL/min). The splitting ratio was 20:1. The methyl esters were positively identified by comparison to known standards (Larodan Fine Chemicals, Malmo, Sweden) and verified by mass spectrometry. Quantification of the fatty acids was made with Chrom-Card A/D 1.0 chromatography station (Carlo Erba Instruments) based on heneicosanoic acid as an internal standard.

Acyl-CoA-esters

Acyl-CoA esters in liver were measured by reversed-phase high-performance liquid chromatography. 100 mg frozen liver was homogenised in ice-cold 1.4 mol/L HClO₄ and 2 mmol/L D-dithiothreitol to obtain 10% (w/v) homogenate, and centrifuged at 12 000 x g for 1 min. 122 µl ice-cold 3 mol/L K₂CO₃ with 0.5 mol/L triethanolamine was added to 500 µl of the supernatant. After 10 min on ice, the solution was centrifuged at 12 000 x g for 1 min at 4°C. 40 µl of the supernatant was injected onto the high-performance liquid chromatography column, and the acyl-CoA esters were measured according to Demoz *et al* (39), with the following modifications: elution buffer A was adjusted to pH 5.00, the profile of the gradient elution was as follows: 0 min, 83.5% A; 10 min, 55% A; 17 min, 10% A, and the flow-rate was 1.0 mL/min.

Isolation of plasma triacylglycerol-rich lipoprotein fraction

Plasma triacylglycerol-rich lipoprotein fraction was prepared by ultracentrifugation of 3 mL plasma at a density of 1.063 g/mL for 19 hr at 105 000 x g at 15°C. The tubes were sliced, and the floating fraction in the top 1 mL of each tube was harvested. The fraction was then dialyzed against 150 mmol/L sodium chloride, 16 mmol/L sodium phosphate and 4 mmol/L potassium phosphate, pH 7.4, saturated with nitrogen.

Real-time quantitative RT-PCR

Total RNA was purified using Trizol (Gibco BRL), and 1 µg total RNA was reversed-transcribed in a total volume of 100 µl by use of a Reverse transcriptase kit (Applied Biosystems). Reactions in which RNA was omitted served as negative control, and reactions in which RNA was diluted served as standard curves.

Primers and Taqman probe for rat Δ^9 , Δ^6 and Δ^5 desaturases, peroxisome proliferator-activated receptor (PPAR) α and glyceraldehyde-3-phosphate dehydrogenase (GAPDH) were designed using Primer Express (Applied Biosystems). GAPDH and 18S rRNA were used as endogenous controls. Primers and Taqman probe for 18S rRNA were purchased from Applied Biosystems.

Real-time PCR was carried out in triplicate for each sample on an ABI 7900 sequence detection system (Applied Biosystems). For Δ^9 , Δ^7 and Δ^5 desaturases, PPAR α and GAPDH, each 20 μ l-reaction contained 3 μ l first-strand cDNA, 1x Universal Master Mix (Applied Biosystems), 300 nmol/L of each forward and reverse primer, and 250 nmol/L Taqman probe. For 18S rRNA the reaction contained 3 μ l first-strand cDNA, 1x Universal Master Mix (Applied Biosystems), and 1x 18S probe/primer reaction mix. All reactions were carried out using the following cycle parameters: 50°C for 2 min and 95°C for 10 min, followed by 40 cycles of 95°C for 15 sec and 60°C for 1 min, as generally recommended by Applied Biosystems. Ct readings (threshold cycle number) for each of the unknown samples were used to calculate the amount of desaturases, PPAR α and GAPDH and 18S rRNA. For each sample, results were normalised to GAPDH and 18S rRNA. Only results normalised to GAPDH are shown, but this was similar to the results normalised to 18 S rRNA.

The results are reported as means \pm SEM from 6 animals in each experimental group. Statistical analysis was by one-way Anova Dunett's test (Prism, GraphPad).

Example 1

Preparation of fish protein hydrolyzate

FPH was produced from fish flesh remnants on salmon bone frames after filleting. Frames without heads from freshly filleted Atlantic salmon (*Salmon salar*, L.) were taken directly from the production line and frozen at -20 ± 2 °C. Within a week the frozen frames were used in the enzymatic hydrolyzing process.

The enzymatic hydrolysis was performed with ProtamexTM at a pH of about 6,5 and at temperature of 55 ± 2 °C. ProtamexTM (E.C. 3.4.21.62/3.4.24.28) is a *Bacillus* protease complex from Novozymes AS (Bagsvaerd, Denmark) and fulfills the purity demands for food-grade enzymes. The ratio of salmon frames to water was 1.14. An enzyme to substrate ratio of 11.1 AU/kg crude protein was used in the hydrolysis. After 60 min of enzymic treatment the temperature was elevated to 98 °C, which was reached after 105 min.

Large bones were retained in the hydrolysing tank, while small bones were removed by filtering the hydrolysate through a mesh. Thereafter the insoluble fraction was removed in a two-phase separator (Westfalia, Germany, SC.35-26-177, 15 kW, 7200 rpm),
5 before the remaining mixture was separated in a three-phase separator (Westfalia, Germany, SB-7-36-+76, 4 kW, 8520 rpm) into salmon oil, emulsion fraction and aqueous fraction. The aqueous fraction fraction was concentrated (NitroAtomizer, Denmark, Falling Film Evaporator, Ff 100), filtered through an ultramembrane with nominal molecular weight limit of 100 000 (PCI membrane systems, UK, PF100, 2,65
10 m²) and finally the ultramembrane filtered fraction (UF fraction) was spray-dried (Niro Atomizer, Denmark, P-63 tower, T_{in} = 200 °C, T_{out} = 84 °C).

The UF fraction is termed fish protein hydrolyzate (FPH) and was used in experiments indicated below. The FPH material contains about 83 % protein, 10 % ash and about 2
15 % lipids, based on dry weight. The amino acid compositions are given in table 2.

Table 2

Total amino acids in the UF fraction obtained by hydrolyses of salmon frames with
Protamex™.

Component	UF fraction
Amino acids (g kg ⁻¹ crude protein)	
Arg	59.4±0.7
His	39±1
Ile	27.5±0.4
Leu	56.4±0.1
Lys	63.7±0.3
Met	22±1
Phe	26.9±0.7
Thr	39±1
Trp	5.3±0.1
Val	35.5±0.4
Ala	74±1
Asn + Asp	73±3
Cysteine (total)	6.1±0.7
Gln + Glu	116±5
Gly	89±3
OH – Pro	20.7±0.7
Pro	47±1
Ser	37±2
Tyr	21±2
Tau	6.2±.1

Example 2.FPH induces a plasma cholesterol lowering effect

5 Obese Zucker rats were offered a diet containing 20% FPH as the sole source of protein. The FPH is a fatty acid free fish protein hydrolysate (FPH), produces as described above.

10 The plasma cholesterol level were reduced by 49% in Zucker rats fed FPH, as compared to rats fed casein as the feed protein. The result is shown in figure 1. The result clearly demonstrates that the FPH decreases the levels of cholesterol in the plasma and can be used as a cholesterol lowering agent.

15 Example 3

FPH decreases the concentration of triacylglycerols in the liver

Figure 2 shows that FPH induces a lowering of the concentration of triacylglycerols (TG) in the liver of about 50%. This indicates that the compound of the present
20 invention can be used as a lipid lowering agent, and for the treatment and prevention of fatty liver.

Example 4FPH inhibits the activity of Acyl-CoA: cholesterol acyltransferase

25

Acyl-CoA:cholesterol acyltransferase (ACAT) catalyses the reaction in which fatty acyl-CoA is esterified to cholesterol. Cholesteryl ester may then be stored in the cytoplasm as lipid droplets or be secreted as part of VLDL together with free cholesterol. Thus, ACAT plays a major role in the VLDL secretion and the subsequent
30 cholesteryl ester accumulation and risk of cardiovascular disease. In the present Zucker rat experiment FPH protein changed the composition of lipid classes in the triacylglycerol-rich lipoprotein fraction, i.e. the cholesteryl ester and phospholipid contents were lower, while the triacylglycerol content was higher than in rats fed casein.

Furthermore, figure 3 shows that the ACAT activity decreased in rats fed FPH protein as compared to those fed casein. As there is strong evidence that increased ACAT activity plays an important role in the progression of atherosclerosis (46-49), this finding indicates that FPH and soy protein are cardioprotective.

5

Figure 3 shows that ACAT activity was reduced about 30% in rats fed FPH as compared to Zucker rats fed casein.

Example 5

10 FPH increases the mitochondrial β -oxidation

Figure 4 shows that FPH increases the mitochondrial β -oxidation. Increased fatty acid oxidation is an important factor behind the lipid lowering effect of FPH. The increased fatty acid catabolism will decrease the amount of fatty acids available for esterification, and thereby reduce the production and secretion of VLDL by the liver. From figure 4 it can be seen that FPH significantly increased the oxidation of palmitoyl-CoenzymeA compared to control.

20 Example 6

FPH interfere with the lipid homeostasis

The present data indicate that fat free FPH material interfere with the lipid homeostasis, and may promote accumulation of endogenous ligands for. In spite of an unchanged hepatic mRNA level of PPAR α (data not shown), the fatty acid composition in liver, plasma and triacylglycerol-rich lipoprotein fraction were changed in rats fed FPH as compared to those fed casein, and the changes did not parallel in liver and plasma (Tables 3-5). The liver concentrations (Table 3) of the saturated 14:0 and 16:0 fatty acids were decreased, while 18:0 was increased in rats fed FPH, in particular those animals fed soy protein, as compared to those fed casein. The liver concentrations of monounsaturated fatty acids, including 18:1n-9, was decreased in rats fed FPH. As a result the 18:1n-9/18:0 ratio was reduced by 54 in rats fed FPH. The hepatic mRNA level of Δ^9 desaturase, however, was not affected by the dietary protein (data not

30

shown). In contrast to liver, an opposite effect was found in plasma on saturated and monounsaturated fatty acids (Table 4). In plasma, the saturated fatty acids 14:0 and 16:0 increased by FPH feeding. The monounsaturated fatty acids 18:1n-9 and 16:1n-7 increased 1.3-1.5 fold in rats fed FPH, and the ratio of 18:1 n-9 to 18:0 increased. In the triacylglycerol-rich lipoprotein fraction (Table 5) there were only minor changes in the saturated and monounsaturated fatty acids, i.e. 16:1n-7 in phospholipids was reduced in rats fed FPH and soy protein. The 18:1n-9/18:0 ratio was generally unchanged in the triacylglycerol-rich lipoprotein fraction. Of the n-6 fatty acids, 18:3n-6 was unchanged in liver, 18:2n-6 increased less than two-fold, while 20:3n-6 and 20:4n-6 increased several-fold with FPH. This resulted in a decrease in the 18:3n-6/18:2n-6 ratio, indicating a reduced Δ^6 desaturation. The hepatic Δ^6 desaturase mRNA expression was indeed reduced (data not shown). Furthermore, the 20:4n-6/20:3n-6 ratio was decreased, indicating a reduced Δ^5 desaturation, in accordance with a decreased hepatic Δ^5 desaturase mRNA level (data not shown). A two-fold increase in the 20:4n-6/18:2n-6 ratio and a 3- and 5-fold increase in the 20:4n-6/18:3n-6 ratio was seen in the animals fed the FPH. It is therefore anticipated that their hepatic elongase activities were increased. Animals fed FPH showed increased plasma concentrations of 18:2 n-6, while they showed decreased plasma concentrations of 20:4 n-6. As a result their 20:4n-6/18:2n-6 ratio in plasma was reduced by 54. Furthermore, the 20:4n-6/20:3n-6 ratio was decreased by 60% in FPH fed rats relative to those fed casein. In addition there was a tendency of decreased 18:3n-6/18:2n-6 and 20:4n-6/18:3n-6 ratios in animals fed FPH, although not significantly. In the triacylglycerol-rich lipoprotein fraction the fatty acid composition resembled the plasma profile, i.e. the 18:2n-6 increased and the 20:4n-6 decreased, resulting in reduced 20:4n-6/18:2n-6 (by 49-68%), 20:4n-6/18:3n-6 (by 53-65%) and 20:4n-6/20:3n-6 (by 35-37%) ratios in rats fed FPH. All of the n-3 fatty acids measured in liver were increased in both FPH fed rats. 18:3n-3 increased 1.8 in plasma by FPH feeding. 20:5n-3 was significantly increased in FPH-fed rats (44% increase), while DHA was reduced by FPH feeding (27 % reduction). In the triacylglycerol-rich lipoprotein fraction the composition of the n-3 fatty acids pattern were changed only in phospholipids, where 20:5n-3 and 22:5n-3 were increased in rats fed FPH. This resembled the findings in plasma.

55215350

20

TABLE 3*Fatty acid composition in liver of Zucker rats fed FPH or casein for 3 weeks¹*

Fatty acids	FPH	Casein
<i>g/100g fat</i>		
14:0	1,5 ± 0,0*	1,7 ± 0,1
16:0	35,2 ± 0,5*	39,3 ± 0,6
18:0	8,8 ± 0,3*	4,9 ± 0,3
Amount saturated	45,5 ± 0,7	45,9 ± 0,6
16:1n-9	0,5 ± 0,0*	0,8 ± 0,0
18:1n-9	24,5 ± 0,7*	29,6 ± 0,4
16:1n-7	6,5 ± 0,3*	8,1 ± 0,5
18:1n-7	2,2 ± 0,2*	3,0 ± 0,1
Amount monounsaturated	33,6 ± 0,9*	41,5 ± 0,8
18:2n-6	11,6 ± 0,1*	8,8 ± 0,5
18:3n-6	0,3 ± 0,0	0,4 ± 0,0
20:3n-6	0,6 ± 0,1*	0,1 ± 0,1
20:4n-6	5,6 ± 0,4*	2,1 ± 0,3
Amount n-6	18,1 ± 0,5*	11,4 ± 0,8
18:3n-3	0,6 ± 0,0*	0,3 ± 0,0
20:5n-3	0,3 ± 0,1	n.d.
22:5n-3	0,4 ± 0,0	n.d.
22:6n-3	1,4 ± 0,2*	0,5 ± 0,1
Amount n-3	2,6 ± 0,2*	0,8 ± 0,1
<i>Ratios</i>		

18:1n-9/18:0	$2,8 \pm 0,2^*$	$6,1 \pm 0,3$
18:3n-6/18:2n-6	$0,028 \pm 0,003^*$	$0,045 \pm 0,003$
20:4n-6/18:2n-6	$0,5 \pm 0,0^*$	$0,2 \pm 0,0$
20:4n-6/18:3n-6	$18,0 \pm 2,5^*$	$5,9 \pm 1,1$

¹ Values are means \pm SEM, n=6.

* different from casein, $P < 0.05$. n.d., not detected.

5

TABLE 4

Fatty acid composition in plasma of Zucker rats fed FPH. or casein for 3 weeks¹

Fatty acids	FPH	Casein
<i>g/100g fat</i>		
14:0	$0,8 \pm 0,1^*$	$0,6 \pm 0,0$
16:0	$21,8 \pm 1,0^*$	$18,6 \pm 0,6$
18:0	$11,8 \pm 0,5$	$12,6 \pm 0,7$
Amount saturated	$33,3 \pm 0,7$	$32,9 \pm 0,8$
16:1n-9	$0,4 \pm 0,0$	$0,4 \pm 0,0$
18:1n-9	$13,6 \pm 0,7^*$	$9,6 \pm 0,5$
16:1n-7	$4,0 \pm 0,3^*$	$3,1 \pm 0,1$
18:1n-7	$1,6 \pm 0,2$	$1,5 \pm 0,1$
Amount monounsaturated	$18,5 \pm 1,4$	$16,1 \pm 1,1$
18:2n-6	$22,0 \pm 0,9^*$	$17,4 \pm 1,0$
18:3n-6	$0,7 \pm 0,1$	$0,6 \pm 0,1$
20:3n-6	$1,3 \pm 0,2$	$0,9 \pm 0,2$

20:4n-6	17,1 ± 1,5*	29,2 ± 1,1
Amount n-6	43,3 ± 1,9	46,3 ± 1,8
18:3n-3	0,8 ± 0,1*	0,5 ± 0,1
20:5n-3	1,0 ± 0,1*	0,6 ± 0,1
22:5n-3	0,6 ± 0,0*	0,4 ± 0,0
22:6n-3	1,6 ± 0,1*	2,3 ± 0,1
Amount n-3	4,0 ± 0,1	3,7 ± 0,1
<i>Ratios</i>		
18:1n-9/18:0	1,2 ± 0,1*	0,8 ± 0,1
18:3n-6/18:2n-6	0,032 ± 0,004	0,037 ± 0,005
20:4n-6/18:2n-6	0,8 ± 0,1*	1,7 ± 0,1
20:4n-6/18:3n-6	35,2 ± 7,4	41,6 ± 6,3
20:4n-6/20:3n-6	14,4 ± 2,2*	37,6 ± 4,6

¹ Values are means ± SEM, n=6.

* different from casein, P < 0.05.

TABLE 5

Fatty acid composition in the different lipid classes in the triacylglycerol-rich lipoprotein fraction of Zucker rats fed FPH or casein for 3 weeks¹

Fatty acids	Triacylglycerol		Phospholipid		Cholesteryl ester	
	FPH	Casein	FPH	Casein	FPH	Casein
<i>g/100g fat</i>						
14:0	1,6 ± 0,1	1,5 ± 0,2	0,5 ± 0,1	0,5 ± 0,0	1,7 ± 0,2	1,2 ± 0,1
16:0	34,9 ± 3,0	33,4 ± 2,8	22,6 ± 0,2	24,7 ± 1,2	19,3 ± 2,1	19,0 ± 0,4
18:0	3,0 ± 0,4	3,2 ± 0,3	29,5 ± 0,7	27,3 ± 1,4	8,2 ± 1,1	3,4 ± 0,4
Σ saturated	39,9 ± 3,4	38,6 ± 3,2	54,8 ± 0,8	54,8 ± 0,3	32,0 ± 3,5	27,7 ± 0,8

55215350

23

16:1n-9	0,5 ± 0,0	0,5 ± 0,1	0,1 ± 0,0	0,1 ± 0,0	0,6 ± 0,1	0,6 ± 0,0
18:1n-9	26,0 ± 0,8	25,5 ± 1,0	5,7 ± 0,5	5,3 ± 0,3	11,8 ± 2,8	12,2 ± 1,8
16:1n-7	6,0 ± 0,5	5,8 ± 0,7	0,7 ± 0,1	0,7 ± 0,0	4,6 ± 1,1	5,8 ± 1,0
18:1n-7	2,6 ± 0,3	3,1 ± 0,2	1,0 ± 0,1*	1,3 ± 0,1	1,0 ± 0,3	1,2 ± 0,3
Σ monounsaturated	35,5 ± 0,8	35,2 ± 1,7	7,9 ± 0,6	8,1 ± 0,2	18,5 ± 4,1	20,0 ± 3,0
18:2n-6	19,4 ± 1,8	19,5 ± 4,3	20,4 ± 0,7	17,5 ± 1,1	17,7 ± 0,6*	14,4 ± 0,2
18:3n-6	0,3 ± 0,0	0,3 ± 0,0	0,1 ± 0,0	0,1 ± 0,0	1,3 ± 0,2	1,0 ± 0,2
20:3n-6	0,4 ± 0,1	0,5 ± 0,1	1,8 ± 0,2	1,2 ± 0,2	1,0 ± 0,1*	0,5 ± 0,1
20:4n-6	1,1 ± 0,1	2,0 ± 0,4	10,8 ± 0,4*	13,8 ± 0,7	27,0 ± 7,3	34,5 ± 3,5
Σ n-6	21,8 ± 2,0	23,0 ± 4,1	33,8 ± 0,8	33,2 ± 0,4	47,0 ± 7,2	50,4 ± 3,8
18:3n-3	1,8 ± 0,0	1,5 ± 0,4	0,1 ± 0,0	0,1 ± 0,0	0,5 ± 0,1	0,5 ± 0,1
20:5n-3	0,4 ± 0,1	0,4 ± 0,1	0,4 ± 0,0*	0,1 ± 0,0	1,2 ± 0,3	0,6 ± 0,2
22:5n-3	0,4 ± 0,0	0,4 ± 0,1	0,7 ± 0,1*	0,5 ± 0,0	n.d.	n.d.
22:6n-3	0,4 ± 0,1	0,6 ± 0,1	2,3 ± 0,4	2,7 ± 0,0	0,7 ± 0,2	0,7 ± 0,0
Σ n-3	2,5 ± 0,8	3,0 ± 0,5	3,4 ± 0,4	3,5 ± 0,0	2,5 ± 0,3	1,8 ± 0,2
Ratios						
18:1n-9/18:0	8,9 ± 1,1	8,0 ± 0,5	0,2 ± 0,0	0,2 ± 0,0	1,4 ± 0,2*	2,2 ± 0,2
18:3n-6/18:2n-6	0,018 ± 0,002	0,017 ± 0,006	0,004 ± 0,002	0,005 ± 0,001	0,071 ± 0,009	0,071 ± 0,014
20:4n-6/18:2n-6	0,058 ± 0,002	0,113 ± 0,032	0,5 ± 0,0*	0,8 ± 0,1	1,5 ± 0,4	2,4 ± 0,2
20:4n-6/18:3n-6	3,3 ± 0,1*	6,9 ± 1,0	92,5 ± 15,6	155,1 ± 19,7	21,2 ± 3,9	36,4 ± 8,7
20:4n-6/20:3n-6	2,8 ± 0,4*	4,3 ± 0,2	6,0 ± 0,8*	12,6 ± 2,1	27,5 ± 6,1*	73,6 ± 11,2

* Values are means ± SEM, n=3 (totally six rats, but plasma from two rats were pooled).

* different from casein, P < 0.05; n.d., not detected; soy, soy protein.

Example 7.

5 FPH lowers the concentration of homocystein in plasma

Increased levels of homocysteine, i.e. hyperhomocysteinemia has been proposed to be associated with arterial diseases, and we thus measured the levels of homocysteine in the plasma samples from rats.

10

Total plasma homocysteine was measured by a fully automated fluorescence assay. 30 µl plasma was reduced by 30 µl NaBH₄/DMSO solution (6 mol/L). After 1,5 min 20 µl of the fluorescence reagent monobromobimane (25 mmol/L) in acetonitrile was added


55215350

24

and allowed to react for 3 min. 20 µl of the sample was then immediately analysed with HPLC by injection on a strong cation-exchange column, and then by column switching into a cyclohexyl silica column. The SCX column was eluted isocratically and the CH column was eluted with a linear methanol gradient (17 - 35% in 5 min) in 20 mmol/L formate buffer. The homocysteine was eluted at a retention time of 4.5 min. The results are given in table 5.

Table 5.Plasma concentration of homocysteine


	Plasma concentration (µmol/L)
Control (casein)	1,37 ± 0,27
FPH	1,17 ± 0,18



CLAIMS

1. Use of a fish protein hydrolysate (FPH) for the preparation of a pharmaceutical or nutritional preparation for the treatment and/or prevention of atherosclerosis,
5 coronary heart disease, stenosis, thrombosis, myocardial infarction, stroke and fatty liver in an animal.
2. Use of a fish protein hydrolysate (FPH) for the preparation of a pharmaceutical or nutritional composition for the treatment and/or prevention of hypercholesterolemia.
10
3. Use of a fish protein hydrolysate (FPH) for the preparation of a pharmaceutical or nutritional composition for lowering the concentration of homocysteine in the plasma.
- 15 4. Use if a fish protein hydrolysate (FPH) for the preparation of a pharmaceutical or nutritional cardio protective composition.
5. Use of a fish protein hydrolysate (FPH) in accordance with one of the preceding claims, wherein said animal is a human.
20
6. Use of a fish protein hydrolysate (FPH) in accordance with one of the preceding claims, wherein said animal is an agricultural animal, such as gallinaceous birds, bovine, ovine, caprine or porcine mammals.
- 25 7. Use of a fish protein hydrolysate (FPH) in accordance with one of the preceding claims, wherein said animal is a domestic or pet animal, such as dog or cat.
8. Use of a fish protein hydrolysate (FPH) in accordance with one of the preceding claims,, wherein said animal is a fish or shellfish, such as salmon, cod, Tilapia, clams,
30 oysters, lobster or crabs.
9. Use in accordance with one of the preceding claims, wherein the fish protein hydrolysate (FPH) is produced by enzymatic hydrolysis of a fish material.

10. Use in accordance with claim 9, wherein the fish material is fish flesh remnants on salmon bone frames after filleting.
- 5 11. Use in accordance with claim 9, wherein the hydrolysis is conducted by the enzyme material ProtamexTM.
12. Use in accordance with one of the claims 1-11, wherein the composition is a food grade product or additive, e.g. an animal feed or pet food.




Abstract

The present invention relates to the use of a fish protein hydrolyzate (FPH). The FPH material lowers the concentration of cholesterol in plasma, and triglycerides in the liver.

- 5 FPH also induces a favourable change in the fatty acid pattern, and lowers the concentration of homocysteine in plasma. A preferable embodiment of the invention relates to the use of FPH as an anti-atherogenic and cardio protective agent, either given as a pharmaceutical or as a functional food.

10

15



55215350

Fig. 1
Cholesterol in plasma






Fig. 2
Triacylglycerols in liver



55215350

Fig. 3







FIG. 4
b-Oxidation in liver M-fraction, palm-carn



b-oxidation in liver M-fraction, palm-CoA



**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.